

Prevalence of patients treated with anti-diabetic medicine in Greenland and Denmark. A cross-sectional register study

Diabetes mellitus is a large and growing worldwide health issue. Prior to this publication, a direct comparison of the prevalence of persons treated with anti-diabetic medicine in Greenland and Denmark has not been found. Therefore, the aim of this study is to estimate and compare the age- and gender-specific prevalence of patients treated with anti-diabetic medicine comparing Greenland and Denmark. The study was performed as a cross-sectional register study using data from population and medical registers in Greenland and Denmark. A total of 784 Greenlandic and 215,580 Danish patients treated with anti-diabetic medicine were included. The prevalence of patients aged 20–79 years treated with anti-diabetic medicine in Greenland was 2.6% (95% CI 2.4–2.8), much lower (p 0.001) compared to Denmark with 5.2% (95% CI 5.2–5.2). The difference was less pronounced after excluding those treated with insulin and women below 45 years treated with metformin. In conclusion, this study showed a lower prevalence of patients treated with anti-diabetic medicine in Greenland than Denmark. The main reason may be a much higher prevalence of undiagnosed diabetes in Greenland, particularly among the middle-aged. Differences in awareness of diabetes and access to continued primary healthcare may be contributing factors.

Forfatter: I Meklenborg ; Michael Lyng Pedersen; Eva Cecilie Bonefeld-Jørgensen **Type:** Article | Artikel
Årstal: 2018 **Emner:** Diabetes mellitus; Prevalence; Greenland; Inuit; Denmark **Titel på tidsskrift:** Journal of Circumpolar Health **Volume på tidsskrift:** 77 **Udgiver:** Taylor & Francis **DOI nummer:**
<https://doi.org/10.1080/22423982.2018.1542930>
